

DTIC FILE COPY

4

OFFICE OF NAVAL RESEARCH

Grant N00014-90-J-1193

TECHNICAL REPORT No. 5

Spontaneous Decay and Resonance Fluorescence of an Admolecule Near
a Silver Surface with Random Roughness

by

Xiao-shen Li, D. L. Lin and Thomas F. George

Prepared for Publication

in

Physical Review B

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

February 1990

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
FEB 16 1990
S D

90 02 15 040

AD-A218 572

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/90/TR-5			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260				7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Grant N00014-90-J-1193	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.		PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Spontaneous Decay and Resonance Fluorescence of an Admolecule Near a Silver Surface with Random Roughness					
12. PERSONAL AUTHOR(S) Xiao-shen Li, D. L. Lin and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) February 1990	
				15. PAGE COUNT 22	
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review B					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	SPONTANEOUS DECAY NEAR SILVER SURFACE		
			RESONANCE FLUORESCENCE RANDOM ROUGHNESS		
			ADMOLECULE THEORY VS EXPERIMENT		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The spontaneous decay rate and resonance fluorescence spectrum of a molecule adsorbed near a rough silver surface are calculated. An assumed random distribution of the roughness on the surface is treated as an effective layer. The calculated decay rate is in excellent agreement with experimental measurements. The effects of surface roughness on both the spontaneous decay rate and resonance fluorescence spectrum are discussed for different cases.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

DD FORM 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

Spontaneous decay and resonance fluorescence of an admolecule
near a silver surface with random roughness

Xiao-shen Li*, D. L. Lin and Thomas F. George
Department of Physics and Astronomy
State University of New York at Buffalo
Buffalo, New York 14260

Abstract

The spontaneous decay rate and resonance fluorescence spectrum of a molecule adsorbed near a rough silver surface are calculated. An assumed random distribution of the roughness on the surface is treated as an effective layer. The calculated decay rate is in excellent agreement with experimental measurements. The effects of surface roughness on both the spontaneous decay rate and resonance fluorescence spectrum are discussed for different cases.

1990 PACS numbers: 32.50.+d, 33.20.Kf, 78.90.+t, 79.20.Ds

* On leave of absence from the Shanghai Institute of Metallurgy, Academia Sinica, Shanghai, People's Republic of China.

Accession For	
NTIS	CRAXI <input checked="" type="checkbox"/>
DTIC	Tab <input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability Codes	
Dist	Avail and/or Special
A-1	

I. Introduction

Fluorescence and light scattering from atoms or molecules near solid surfaces have aroused much interest since the perfection of the fatty-acid monolayer assembly technique¹ has led to a series of successful measurements of the lifetime of excited molecules near metal surfaces.² Despite other possible effects,³ the role of surface roughness has been recognized to be prominent in various surface processes⁴ ever since the discovery of the surface-enhanced Raman effect⁵ in 1974. Processes such as photolysis and photochemical degradation have been studied in great detail by theoretical models including both localized⁶ and extended⁷ surface structures. It has become clear that the ultimate outcome for the enhancement of photoabsorption and resonance fluorescence processes generally depends on two competing factors. One is the enhanced surface electromagnetic field, and the other is the surface-induced decay rate of adatoms.^{4,8}

Effects of the surface roughness on the spontaneous decay rate of adsorbed molecules has been investigated both theoretically⁹ and experimentally.¹⁰ It is found that the decay rate increases due to the presence of surface roughness. As a matter of fact, the decay rate of pyrazine molecules near a silver surface is measured to be about five times larger than the theoretically expected value which is calculated for a flat smooth surface. Such a tremendous increase, as has been suggested recently,¹¹ can be attributed to the roughness of the silver surface by assuming periodically roughened metallic surfaces.

Resonance fluorescence of adatoms at rough metallic surfaces, on the other hand, has also been studied¹² by considering the roughness as hemispherical protrusions on perfectly conducting solid surfaces. We consider, in this article, the spontaneous decay rate and resonance

fluorescence spectrum of a molecule adsorbed near a rough silver surface. The roughness is assumed to be small clusters of atoms, that may or may not be silver, distributed randomly on the surface, and is treated as an effective layer¹³ with its effective optical constants modeled by the Maxwell-Garnett theory.¹⁴ This treatment has been proven very successful in the investigation of optical excitation of surface plasma waves along rough silver surfaces. Our results are found to be in excellent agreement with experiments.

II. Theory

Consider a molecule with two levels $|+\rangle$ and $|-\rangle$ separated by a distance $\hbar\omega$. It is adsorbed near a bulk silver with rough surface. The molecule is driven by a monochromatic laser field with

$$E(t) = \frac{1}{2} (E e^{i\omega_0 t} + E^* e^{-i\omega_0 t}) \quad (1)$$

We can apply the surface-dressed optical Bloch equation^{15,16} (SBE) to describe the interaction process. The SBE are given by¹⁶

$$\frac{d}{dt} \begin{bmatrix} \langle S^+ \rangle \\ \langle S^z \rangle \\ \langle S^- \rangle \end{bmatrix} = \begin{bmatrix} i(\Delta + \Omega^S) - \gamma & i\Omega & 0 \\ i\Omega^*/2 & -2\gamma & -i\Omega/2 \\ 0 & -i\Omega^* & -i(\Delta + \Omega^S) - \gamma \end{bmatrix} \begin{bmatrix} \langle S^+ \rangle \\ \langle S^z \rangle \\ \langle S^- \rangle \end{bmatrix} - \begin{bmatrix} 0 \\ \gamma \\ 0 \end{bmatrix} \quad (2)$$

The notation is as follows. The adatom with a transition frequency ω is located at a distance d away from the surface of silver. The matrix element of the electric dipole moment operator is denoted by $|p|$, and E and ω_0 are the amplitude and frequency of the external laser field, respectively. The

detuning is $\Delta = \omega - \omega_0$, and the Rabi frequency is $\Omega = |p|E$. The transition probability amplitude is proportional to the projection operators defined by

$$\begin{aligned} S^+ &= |+\times-| \\ S^z &= \frac{1}{2} (|+\times+| - |-\times-|) \\ S^- &= |-\times+| \end{aligned} \quad (3)$$

The total decay rate of the adatom can be written as

$$\gamma = \gamma^0 + \gamma^s, \quad (4)$$

where γ^0 is the decay rate in the absence of the substrate,

$$\gamma^0 = \frac{2}{3} \sqrt{\epsilon} |p|^2 \omega^3 / c^3, \quad (4a)$$

and

$$\gamma^s = |p|^2 \operatorname{Im} f(d) \quad (4b)$$

is the decay rate induced by the surface. The frequency shift of the spontaneous radiation due to the surface is

$$\Omega^s = |p|^2 \operatorname{Re} f(d), \quad (5)$$

and the function $f(d)$ is introduced only for convenience¹² and is determined by¹⁶

$$E_R = |p| f(d) S^- = pf(d) \quad (6)$$

where E_R is the component of the reflected field \vec{E}_R in the direction of \vec{p} .

Equation (2) agrees with the result from linear response theory¹⁷ when the adatom is taken as a harmonic oscillator. When the intensity of the incident laser is weak, it is more probable to find the molecule in its lower state. Thus we can take $\langle S^z \rangle \approx -\frac{1}{2}$. The SBE (2) can then be linearized and solved. With the initial condition¹⁶ $\langle S^2(0) \rangle = -\frac{1}{2}$, we find the solution

$$\langle S^2(t) \rangle = |\Omega|^2 [1 + e^{-2\gamma t} - 2e^{-\gamma t} \cos(\Delta + \Omega^S)t] / 4|z|^2 - \frac{1}{2}, \quad (7)$$

where $z = \gamma + i(\Delta + \Omega^S)$. By making use of the regression theorem for correlation functions,¹⁸ we find from the SBE the well-known results of the incoherent resonance fluorescence spectrum,^{18,19}

$$\bar{g}(\nu) = \frac{1}{2} |\Omega|^4 \gamma (D^2 + \frac{1}{2} |\Omega|^2 + 4\gamma^2) / (\frac{1}{2} |\Omega|^2 + |z|^2) (x^2 + y^2), \quad (8)$$

where $D = \nu - \omega_0$, $x = 2\gamma(\frac{1}{2} |\Omega|^2 + |z|^2 - 2D^2)$, and $y = D(|\Omega|^2 + |z|^2 + 4\gamma^2 - D^2)$.

It is seen from Eqs. (4)-(8) that both the spontaneous emission properties and the resonance fluorescence spectrum depend sensitively on the surface-reflected field \vec{E}_R which, in turn, depends on the optical constant of silver and the structure of the silver surface. In these equations, we have regarded the molecule as an emitting dipole. The emitted field is reflected back from the surface and is coupled to the radiation dipole whose dynamical behavior is therefore totally changed. To calculate the reflected field, we must take into account the roughness of the silver surface. This is treated

in a similar fashion as is discussed in Ref. 13. The roughness is assumed to be on a very fine scale which is much smaller than the wavelength of the relevant light. The scattering of light due to the roughness is then very small and consequently the roughness can be modeled by introducing an effective layer on the silver surface. The optical constant ϵ_2 of this layer is evaluated by the Maxwell-Garnett theory

$$\epsilon_e = \epsilon [\epsilon_s(1+2q) + 2\epsilon(1-q)] / [\epsilon_s(1-q) + \epsilon(2+q)], \quad (9)$$

where ϵ is the dielectric constant of the medium containing the molecule, ϵ_s stands for the complex dielectric constant of the silver substrate, and q represents the volume fraction of silver in the effective thin layer. It is noted that the maximum roughness corresponds to $q = 0.5$.

The reflected electric field at the dipole can be found by classical electromagnetic field theory.^{16,20} The results are, with $x_0 = \omega/c$,

$$E_R^{\parallel} = \sqrt{\epsilon} \kappa_0^3 p \int_0^{\infty} d\kappa \kappa \mu^{-1} (R_{\perp} + \mu^2 R_{\parallel}) \quad (10)$$

when the dipole moment \vec{p} is parallel to the surface and

$$E_R^{\perp} = -i\sqrt{\epsilon} \kappa_0^3 p \int_0^{\infty} d\kappa \kappa^3 \mu^{-1} R_{\parallel} \quad (11)$$

when \vec{p} is normal to the surface. Here we have introduced $\mu = \sqrt{1 - \kappa^2}$ and

$$R_{\perp, \parallel} = \frac{R_{\perp, \parallel}^m + R_{\perp, \parallel}^e}{1 + R_{\perp, \parallel}^m R_{\perp, \parallel}^e} e^{i2\mu d}$$

with

$$R_{\parallel}^m = \frac{\epsilon \mu_e - \epsilon_e \mu}{\epsilon \mu_e + \epsilon_e \mu} \quad (13a)$$

$$R_{\perp}^m = \frac{\mu - \mu_e}{\mu + \mu_e} \quad (13b)$$

$$R_{\parallel}^e = \frac{\epsilon_e \mu_s - \epsilon_s \mu_e}{\epsilon_e \mu_s + \epsilon_s \mu_e} e^{i2\mu_e \hat{d}_e} \quad (14a)$$

$$R_{\perp}^e = \frac{\mu_e - \mu_s}{\mu_e + \mu_s} e^{i2\mu_e \hat{d}_e} \quad (14b)$$

$$\mu_e = \sqrt{\epsilon_e / \epsilon - \kappa^2} \quad (15a)$$

$$\mu_s = \sqrt{\epsilon_s / \epsilon - \kappa^2} \quad (15b)$$

$$\hat{d} = 2\pi\sqrt{\epsilon}d/\lambda \quad (16a)$$

and the thickness of the effective layer

$$\hat{d}_e = 2\pi\sqrt{\epsilon}d_e/\lambda \quad (16b)$$

Substituting (10) and (11) in (6), we can rewrite γ and Ω^S in (4) and (5) for dipole orientations parallel and normal to the surface,

$$\gamma_{\parallel} = 1 + \frac{3}{4} \operatorname{Re} \int_0^{\infty} d\kappa \kappa (R_{\perp} + \mu^2 R_{\parallel}) / \mu \quad (17a)$$

$$\gamma_{\perp} = 1 - \frac{3}{2} \operatorname{Re} \int_0^{\infty} \kappa^3 d\kappa R_{\perp} / \mu \quad (17b)$$

$$\Omega_{\parallel}^S = - \frac{3}{4} \operatorname{Im} \int_0^{\infty} \kappa d\kappa (R_{\perp} + \mu^2 R_{\parallel}) / \mu \quad (18a)$$

$$\Omega_{\perp}^S = \frac{3}{4} \operatorname{Im} \int_0^{\infty} \kappa^3 d\kappa R_{\parallel} / \mu \quad (18b)$$

where we have taken the unit $\gamma^0 = 1$.

III. Results and discussion

The integrals involved in (17) and (18) can only be evaluated numerically. We choose $\epsilon = 1.7$, $q = 0.2$ and $d_e = 10 \text{ \AA}$ in our calculation. In the neighborhood of the plasmon resonance, we consider two wavelengths, $\lambda = 3800 \text{ \AA}$ and 3200 \AA , for transitions of the adsorbed molecule. The dielectric constants of silver corresponding to these wavelengths are^{10,21} $\epsilon_s(\lambda = 3800) = -3.16 + 0.29i$ and $\epsilon_s(\lambda = 3200) = 0.50 + 0.04i$, respectively. The roughness effect on the wavelength dependence of the decay rate is also investigated for the whole wavelength range.

We first consider the decay rate of the adsorbed molecule as a function of its distance d from the surface. In Figs. 1 and 2, the results computed from (17) are plotted for the two wavelengths. It is clearly observed that for $\lambda = 3800 \text{ \AA}$, the surface roughness enhances the decay rate whether the dipole orientation is parallel or normal to the surface. On the contrary, the roughness prolongs the lifetime of the excited molecule for $\lambda = 3200 \text{ \AA}$. Similar conclusion has been reported in Ref. 11 in which the more special case of periodically roughened metallic surface is considered. Our results are

shown in Fig. 3 in excellent agreement with experimental data.¹⁰ The dashed line representing results calculated for a smooth surface is also plotted for comparison purposes. Therefore, surface roughness plays a very important role indeed and must be included in the treatment of any optical problem involving absorption.

To find how the effect of the roughness on the decay rate varies with the wavelength, we have computed the ratio R of the decay rate for a rough silver surface to that for a flat one as a function of the wavelength when the molecule is located very close to the surface. It is found that in the neighborhood of the surface plasmon resonance, the energy transfer from the adatom to the silver substrate is greatly enhanced due to the roughness. The results are depicted in Fig. 4. It is clearly seen that the roughness has very little effect on the decay rate for long wavelengths, say, $\lambda \geq 6000 \text{ \AA}$. In the shorter wavelength region, say, $\lambda \leq 3300 \text{ \AA}$, the effective layer of roughness that absorbs less energy plays a relatively important role in reducing the energy transfer. Such phenomena can be understood in the following manner. In the region where the radiation wavelengths are far away from the plasmon resonance, the energy transfer process depends mainly upon the imaginary part of the dielectric constant of the substrate. In other words, it depends on the energy absorption rate. For silver, $\text{Im}\epsilon_s$ becomes smaller as the wavelength becomes longer. Since the effective layer of surface roughness has a screening effect on the energy transfer, stronger screening is expected when $\text{Im}\epsilon_s$ is larger or when the wavelength is shorter.

We have also investigated the variation of the roughness effect with the distance by calculating the ratio R as a function of the distance d . As expected, this effect increases as d decreases. Thus, whether the surface roughness enhances or suppresses the spontaneous emission rate depends on both

λ and d . Figure 5 shows a typical case in which we have chosen a wavelength around the surface plasmon resonance in the short wavelength region. It is observed that the "screening effect" of the roughness layer dominates ($R < 1$) for small d , and $R > 1$ for large d . Although the results shown in Figs. 4 and 5 are calculated for the normal orientation of the adatomic dipole moment, we have obtained similar conclusions for the parallel orientation.

We now turn our attention to the interaction of an external driving field with the molecule. In what follows, we only consider the parallel orientation of the dipole moment for simplicity. The normal orientation can be treated in the same fashion, and no qualitatively different conclusion is expected. As we have noted above, we use γ^0 as the unit for all quantities with the dimension t^{-1} . The behavior of the mean molecular inversion as a function of time is calculated from (7) and is shown in Fig. 6. Strong dependence of the results upon the wavelength is obvious as the only difference in (a) and (b) is in the wavelength. This qualitatively different appearance of the curves is primarily due to the energy transfer from the excited molecule to the silver surface. In the case (a), such transfer is weak and the Rabi oscillation of $S^Z(t)$ is evident. The surface roughness has less effect in this case. The situation is totally different in case (b). The energy transfer is so strong that Rabi oscillation can hardly occur and the system goes into steady-state right after the interaction starts. It is also observed that the influence of the surface roughness is very important in this case.

As to the incoherent part of the resonance fluorescence spectrum, we report in Fig. 7 some of our results computed from (8) for (a) $\lambda = 3200 \text{ \AA}$ and (b) $\lambda = 3800 \text{ \AA}$. Once again, the influence of surface roughness depends strongly upon the wavelength. While the two sidebands of the spectrum appear

to be completely suppressed in (b) by the surface roughness, the spectrum does not show much quantitative changes due to the roughness in (a).

Acknowledgments

Mr. Yue Liu has helped with part of the numerical calculations in this work and is gratefully acknowledged. This work was supported in part by the Office of Naval Research and the National Science Foundation under Grant CHE-8620274.

References

1. H. Kuhn, Pure Appl. Chem. 11, 345 (1965); Naturwissenschaften 54, 429 (1967).
2. K. H. Drexhage, H. Kuhn and F. P. Schofer, Ber. Bunsenges. Phys. Chem. 72, 329 (1968); K. H. Drexhage, Sci. Am. 222, 108 (1970).
3. See, e.g., M. Moskovits, Rev. Mod. Phys. 57, 783 (1985).
4. For reviews, see G. M. Goncher, C. A. Parsons and C. B. Harris, J. Phys. Chem. 88, 4200 (1984); D. A. Jelski, P. T. Leung and T. F. George, Int. Rev. Phys. Chem. 7, 179 (1988).
5. M. Fleischmann, P. J. Handra and A. J. McQuillan, Chem. Phys. Lett. 26, 163 (1974).
6. J. I. Gersten and A. Nitzan, Surf. Sci. 158, 165 (1985), and references therein.
7. P. T. Leung and T. F. George, J. Chem. Phys. 85, 4729 (1986); P. T. Leung and T. F. George, Chem. Phys. Lett. 134, 375 (1987).
8. P. T. Leung and T. F. George, Spectroscopy 4, 35 (1989).
9. J. Arias, P. K. Aravind and H. Metiu, Chem. Phys. Lett. 85, 404 (1982).
10. R. Rossetti and L. E. Brus, J. Chem. Phys. 76, 1146 (1982).
11. P. T. Leung, Z. C. Wu, D. A. Jelski and T. F. George, Phys. Rev. B 36, 1475 (1987).
12. X. Y. Huang, K. T. Lee and T. F. George, J. Chem. Phys. 85, 567 (1986); J. T. Lin, X. Y. Huang and T. F. George, J. Opt. Soc. AM. B 4, 219 (1987).
13. G. J. Kovacs and G. D. Scott, Phys. Rev. B 16, 1297 (1977).
14. J. C. Maxwell-Garnett, Philos. Trans. R. Soc. Lond. A 203, 385 (1904); 205, 237 (1906).

15. X. Y. Huang, J. T. Lin and T. F. George, J. Chem. Phys. 80, 893 (1984);
X. Y. Huang and T. F. George, J. Phys. Chem. 88, 4801 (1984); H. F.
Arnoldus, S. van Smaalen and T. F. George, Adv. Chem. Phys. 73, 679
(1989).
16. X. S. Li and C. D. Gong, Phys. Rev. A 35, 1595 (1987); Phys. Rev. B 39,
8284 (1989); B 39, 8293 (1989).
17. G. S. Agarwal, Phys. Rev. A 12, 1475 (1975).
18. M. Lax, Phys. Rev. 157, 213 (1967).
19. B. R. Mollow, Phys. Rev. 188, 1969 (1969); Phys. Rev. A 15, 1023 (1977).
20. R. R. Chance, A. Prock and R. Silbey, Adv. Chem. Phys. 37, 1 (1978).
21. P. B. Johnson and R. W. Christy, Phys. Rev. B 6 4370 (1972).

Figure captions

1. $\log \gamma$ as a function of $\log d$ for $\lambda = 3800 \text{ \AA}$ for the cases of a rough surface (solid line) and smooth surface (dashed line). The dipole moment is oriented (a) normal to the surface and (b) parallel to the surface.
2. Same as in Fig. 1 except that $\lambda = 3200 \text{ \AA}$.
3. A comparison of theoretical and experimental results for the case of parallel dipole orientation. The dashed line representing smooth-surface results is plotted only to show the importance of the roughness effect.
4. The ratio R of decay rates as a function of the wavelength λ for the silver substrate. The distance is $d = 20 \text{ \AA}$. The horizontal line represents $R = 1$.
5. R as a function of d for silver substrate with $\lambda = 3317 \text{ \AA}$. The horizontal line represents $R = 1$.
6. Time evolution of the quantity $(\langle S^z \rangle + \frac{1}{2})/|\Omega|^2$ for $d = 125 \text{ \AA}$ and $\Delta = 0$. The dipole moment is parallel to the surface. The solid (dashed) line represents results calculated for the rough (smooth) surface. (a) $\lambda = 3200 \text{ \AA}$, (b) $\lambda = 3800 \text{ \AA}$.
7. Resonance fluorescence spectrum for the dipole moment parallel to the surface and $|\Omega| = 50$, $\Delta = 0$, $d = 125 \text{ \AA}$. The solid (dashed) line represents rough (smooth) surface results. (a) $\lambda = 3200 \text{ \AA}$, (b) $\lambda = 3800 \text{ \AA}$.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6







Fig. 7



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 M. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853